Table 175. Reported AIDS Cases for Adults and Adolescents by Transmission Category and Sex: 2005

[Provisional. For cases reported in the year shown. Includes Puerto Rico, Virgin Islands, Guam, and U.S. Pacific Islands. Acquired immunodeficiency syndrome (AIDS) is a specific group of diseases or conditions which are indicative of severe immunosuppression related to infection with the human immunodeficiency virus (HIV). Data are subject to retrospective changes and may differ from those data in Table 173]

Transmission category	2005			Cumulative through 2005 ¹		
	Total	Male	Female	Total	Male	Female
Persons 13 years old and over, total Male-to-male sexual contact	41,900 14,819 6,215	30,956 14,819 4,168	10,944 (X) 2,047	946,578 416,232 225,210	764,808 416,232 159,676	181,769 (X) 65,534
injection drug use Hemophilia/coagulation disorder High-risk heterosexual contact Sex with injection drug user Sex with bisexual male Sex with person with hemophilia Sex with HIV-infected transfusion recipient. Sex with HIV-infected person,	1,742 79 8,028 1,140 223 16 40	1,742 68 3,110 389 (X) 3 13	(X) 11 4,918 751 223 13 27	62,940 5,482 126,405 36,681 4,876 574 1,296	62,940 5,142 46,533 11,371 (X) 81 529	(X) 340 79,872 25,310 4,876 493 767
risk factor not specified	6,609	2,705	3,904	82,978	34,552	48,426
components, or tissue	160 10,857	73 6,976	87 3,881	9,334 100,975	5,190 69,095	4,144 31,879

X Not applicable.

Includes persons with a diagnosis of AIDS, reported from the beginning of the epidemic through 2005. Cumulative total includes persons with characteristics unknown.

Source: U.S. Centers for Disease Control and Prevention, Atlanta, GA, HIV/AIDS Surveillance Report, 2005, Volume 17. http://www.cdc.gov/hiv/topics/surveillance/resources/reports/2005report/default.htm (accessed 20 December 2006).